

Contents

Introduction	1
Feedstuff/Diet Evaluation	
Use of defatted soybean meal as a substitute for fish meal in diets of Japanese flounder (<i>Paralichthys olivaceus</i>) K. Kikuchi (Abiko, Japan)	3
High inclusion levels of poultry meals and related byproducts in diets for gilthead seabream <i>Sparus aurata</i> L. I. Nengas, M.N. Alexis (Athens, Greece) and S.J. Davies (Plymouth, UK)	13
Growth of mediterranean yellowtail (<i>Seriola dumerilii</i>) fed extruded diets containing different levels of protein and lipid M. Jover (Valencia, Spain), A. García-Gómez (Murcia, Spain), A. Tomás (Valencia, Spain), F. de la Gándara (Murcia, Spain) and L. Pérez (Valencia, Spain)	25
Growth, feed utilization and body lipid content of gilthead seabream (<i>Sparus aurata</i>) fed increasing lipid levels and fish meals of different quality J.M. Vergara, G. López-Calero, L. Robaina, M.J. Caballero, D. Montero, M.S. Izquierdo (Las Palmas de Gran Canaria, Spain) and A. Aksnes (Bergen, Norway)	35
Digestibility, postprandial ammonia excretion and selected plasma metabolites in European sea bass (<i>Dicentrarchus labrax</i>) fed pelleted or extruded diets with or without wheat gluten L. Robaina (Las Palmas de Gran Canaria, Spain), G. Corraze, P. Aguirre, D. Blanc (Saint Peé sur Nivelle, France), J.P. Melcion (Nantes, France) and S. Kaushik (Saint Peé sur Nivelle, France)	45
Alternative methods for nutrition research on the southern bluefin tuna, <i>Thunnus maccoyii</i> : in vitro digestibility C.G. Carter, M.P. Bransden (Launceston, Tas., Sydney, NSW, Australia), R.J. van Barneveld and S.M. Clarke (Sydney, NSW, Adelaide, SA, Australia)	57
Re-evaluation of crude fibre and acid-insoluble ash as inert markers, alternative to chromic oxide, in digestibility studies with rainbow trout (<i>Oncorhynchus mykiss</i>) A.E. Morales, G. Cardenete, A. Sanz and M. de la Higuera (Granada, Spain)	71
Determination of digestibility of commercial salmon feeds M. Hillestad, T. Asgård and G.M. Berge (Sunndalsøra, Norway)	81
Maintenance requirement for digestible energy and efficiency of utilisation of digestible energy for retention in rainbow trout, <i>Oncorhynchus mykiss</i> M. Rodehutscord and E. Pfeffer (Bonn, Germany)	95
Potential for dietary phytase to improve the nutritive value of canola protein concentrate and decrease phosphorus output in rainbow trout (<i>Oncorhynchus mykiss</i>) held in 11°C fresh water I. Forster, D.A. Higgs (Vancouver, BC, Canada), B.S. Dosanjh (Surrey, BC, Canada), M. Rowshandeli (Vancouver, BC, Canada) and J. Parr (Georgetown, ON, Canada)	109
Phosphorus utilization by rainbow trout (<i>Oncorhynchus mykiss</i>): estimation of dissolved phosphorus waste output D.P. Bureau and C.Y. Cho (Guelph, Ont., Canada)	127

Effect of variations in daily feeding frequency and ration size on growth of shrimp, <i>Litopenaeus vannamei</i> (Boone), in zero-water exchange culture tanks M. Velasco, A.L. Lawrence and F.L. Castille (Port Aransas, TX, USA)	141
Proteins and Amino Acids	
Alternative dietary protein sources for farmed tilapia, <i>Oreochromis</i> spp. A.-F.M. El-Sayed (Alexandria, Egypt)	149
Quantitative dietary requirements of postlarval tiger shrimp, <i>Penaeus monodon</i> , for histidine, isoleucine, leucine, phenylalanine and tryptophan O.M. Millamena, M.B. Teruel (Iloilo, Philippines), A. Kanazawa and S. Teshima (Kagoshima, Japan)	169
In vitro uptake and interaction between arginine and lysine in the intestine of Atlantic salmon (<i>Salmo salar</i>) G.E. Berge (Bergen, Norway), A.M. Bakke-McKellep (Oslo, Norway) and E. Lied (Bergen, Norway)	181
The utilization of diets containing high levels of fish protein hydrolysate by turbot (<i>Scophthalmus maximus</i>) juveniles A. Oliva-Teles, A.L. Cerqueira and P. Gonçalves (Porto, Portugal)	195
Liver and white muscle protein turnover rates in the European eel (<i>Anguilla anguilla</i>): effects of dietary protein quality M. de la Higuera, H. Akharbach, M.C. Hidalgo (Granada, Spain), J. Peragón (Jaén, Spain), J.A. Lupiáñez and M. García-Gallego (Granada, Spain)	203
Lipids and Fatty Acids	
Lipid nutrition of marine fish during early development: current status and future directions J. Sargent, L. McEvoy, A. Estevez, G. Bell, M. Bell, J. Henderson and D. Tocher (Stirling, UK)	217
Effects of 1,2-di-20:5-phosphatidylcholine (PC) and 1,2-di-22:6-PC on growth and stress tolerance of Japanese flounder (<i>Paralichthys olivaceus</i>) larvae A. Tago, Y. Yamamoto, S.-i. Teshima and A. Kanazawa (Kagoshima, Japan)	231
Effect of coconut oil and tricaprylin vs. triolein on survival, growth and fatty acid composition of common carp (<i>Cyprinus carpio</i> L.) larvae S. Fontagné (Saint-Pée-sur-Nivelle, France), T. Pruszyński (Chybie, Poland), G. Corraze and P. Bergot (Saint-Pée-sur-Nivelle, France)	241
Effect of different dietary polar lipid levels and different $n-3$ HUFA content in polar lipids on gut and liver histological structure of gilthead seabream (<i>Sparus aurata</i>) larvae M. Salhi, C.M. Hernández-Cruz (Las Palmas, Spain), M. Bessonart (Montevideo, Uruguay), M.S. Izquierdo and H. Fernández-Palacios (Las Palmas, Spain)	253
Effect of dietary arachidonic acid levels on growth and survival of gilthead sea bream (<i>Sparus aurata</i> L.) larvae M. Bessonart (Montevideo, Uruguay), M.S. Izquierdo (Las Palmas, Spain), M. Salhi (Telde, Spain), C.M. Hernández-Cruz (Las Palmas, Spain), M.M. González and H. Fernández-Palacios (Telde, Spain)	265
Combined effect of lipid level and fish meal quality on liver histology of gilthead seabream (<i>Sparus aurata</i>) M.J. Caballero, G. López-Calero, J. Socorro, F.J. Roo, M.S. Izquierdo and A.J. Fernández (Las Palmas de Gran Canaria, Spain)	277
Study of the $n-3$ highly unsaturated fatty acids requirement and antioxidant status of <i>Dentex dentex</i> larvae at the <i>Artemia</i> feeding stage G. Mourente (Cádiz, Spain), D.R. Tocher (Stirling, UK), E. Diaz-Salvago (Cádiz, Spain), A. Grau and E. Pastor (Majorca, Spain)	291
Relationships between antioxidants, antioxidant enzyme activities and lipid peroxidation products during early development in <i>Dentex dentex</i> eggs and larvae G. Mourente (Cadiz, Spain), D.R. Tocher (Stirling, UK), E. Diaz (Cadiz, Spain), A. Grau and E. Pastor (Palma de Mallorca, Spain)	309

Effect of dietary lipid level on growth performance and feed utilization by European sea bass juveniles (<i>Dicentrarchus labrax</i>) H. Peres and A. Oliva-Teles (Porto, Portugal)	325
Dietary lipid composition affects blood leucocyte fatty acid compositions and plasma eicosanoid concentrations in European sea bass (<i>Dicentrarchus labrax</i>) B.M. Farndale, J.G. Bell, M.P. Bruce, N.R. Bromage (Stirling, UK), F. Oyen, S. Zanuy (Castellón, Spain) and J.R. Sargent (Stirling, UK)	335
The effects of dietary fat and NFE levels on growing European sea bass (<i>Dicentrarchus labrax</i> L.). Growth rate, body and fillet composition, carcass traits and nutrient retention efficiency D. Lanari (Udine, Italy), B.M. Poli (Firenze, Italy), R. Ballestrazzi (Udine, Italy), P. Lupi (Firenze, Italy), E. D'Agaro (Udine, Italy) and M. Mecatti (Firenze, Italy)	351
Metabolism of ursodeoxycholic acid in the Japanese flounder (<i>Paralichthys olivaceus</i>) and the yellowtail (<i>Seriola quinqueradiata</i>) S.-i. Teshima, M. Ishikawa, S. Koshio, M. Yunoki, A. Kanazawa (Kagoshima, Japan) and S. Hayashida (Fukuoka, Japan)	365
Kinetic behavior and protein expression of hepatic NADPH-production systems during development of rainbow trout (<i>Oncorhynchus mykiss</i>) J.B. Barroso, J. Peragón (Jaén, Spain), L. García-Salguero, M. de la Higuera and J.A. Lupiáñez (Granada, Spain)	375
Performance of <i>Macrobrachium rosenbergii</i> broodstock fed diets with different fatty acid composition R.O. Cavalli, P. Lavens and P. Sorgeloos (Ghent, Belgium)	387
Butylated hydroxytoluene: its effect on the quality of shrimp diet stored at various temperatures and on growth and survival of <i>Penaeus monodon</i> juveniles M.N. Bautista-Teruel and P.F. Subosa (Iloilo, Philippines)	403
Carbohydrates	
Glucose tolerance in gilthead seabream (<i>Sparus aurata</i>) and European seabass (<i>Dicentrarchus labrax</i>) H. Peres, P. Gonçalves and A. Oliva-Teles (Porto, Portugal)	415
Carbohydrates affect protein-turnover rates, growth, and nucleic acid content in the white muscle of rainbow trout (<i>Oncorhynchus mykiss</i>) J. Peragón, J.B. Barroso (Jaén, Spain), L. García-Salguero, M. de la Higuera and J.A. Lupiáñez (Granada, Spain)	425
Dietary supplementation of chitin and chitosan depresses growth in tilapia, <i>Oreochromis niloticus</i> × <i>O. aureus</i> S.-Y. Shiau and Y.-P. Yu (Keelung, Taiwan)	439
Vitamin C	
Tissue ascorbic acid levels in European sea bass (<i>Dicentrarchus labrax</i>) and gilthead sea bream (<i>Sparus aurata</i> L.) fingerlings fed diets containing different forms of ascorbic acid M.N. Alexis, I. Nengas, E. Fountoulaki, E. Papoutsi, A. Andriopoulou, M. Koutsodimou (Athens, Greece) and J. Gaubaudan (Saint-Louis, France)	447
Influence of dietary ascorbate derivatives on tissue copper, iron and zinc concentrations in grass shrimp, <i>Penaeus monodon</i> T.-S. Hsu and S.-Y. Shiau (Keelung, Taiwan)	457
Larval Nutrition	
Development of digestive enzymes in larvae of <i>Solea senegalensis</i> , Kaup 1858 L. Ribeiro (Faro, Portugal), J.L. Zambonino-Infante, C. Cahu (Plouzané, France) and M.T. Dinis (Faro, Portugal)	465
Influence of size and nutritional value of <i>Artemia franciscana</i> on growth and quality of halibut larvae (<i>Hippoglossus hippoglossus</i>) during the live feed period A.I. Olsen (Trondheim, Norway), Y. Attramadal (Øyestranda, Norway), A. Jensen and Y. Olsen (Trondheim, Norway)	475

Rearing techniques for red porgy (<i>Pagrus pagrus</i>) during larval development C.M. Hernández-Cruz (Las Palmas de Gran Canaria, Spain), M. Salhi (Telde, Spain), M. Bessonart (Montevideo, Uruguay), M.S. Izquierdo (Las Palmas de Gran Canaria, Spain), M.M. González and H. Fernández-Palacios (Telde, Spain)	489
Development of red porgy <i>Pagrus pagrus</i> visual system in relation with changes in the digestive tract and larval feeding habits F.J. Roo, J. Socorro, M.S. Izquierdo, M.J. Caballero, C.M. Hernández-Cruz, A. Fernández and H. Fernández-Palacios (Las Palmas, Spain)	499
Nutrition and Health	
Nitric oxide production in activated macrophages from channel catfish (<i>Ictalurus punctatus</i>): influence of dietary arginine and culture media J.A. Buentello and D.M. Gatlin III (College Station, TX, USA)	513
Dietary and culture influences on macrophage aggregate parameters in gilthead seabream (<i>Sparus aurata</i>) juveniles D. Montero (Las Palmas, Spain), V.S. Blazer (Kearneysville, WV, USA), J. Socorro, M.S. Izquierdo (Las Palmas de Gran Canaria, Spain) and L. Tort (Barcelona, Spain)	523

